

## Article

## Development of the First Medicinal Crop-Varieties with Enhanced Bioactive Compounds from Bitter Gourd (*Momordica charantia* L.)

Chittaranjan Kole<sup>1,2,\*</sup>, Phullara Kole<sup>1,2</sup>

<sup>1</sup> Department of Genetics and Biochemistry, Clemson University, Clemson, SC 29634, USA; pkole62@gmail.com (PK)

<sup>2</sup> Prof. Chittaranjan Kole Foundation for Science & Society, 1399 Panchpota, Garia, Kolkata 700152, West Bengal, India

\* Correspondence: Chittaranjan Kole, Email: ckoleorg@gmail.com.

---

### ABSTRACT

Incidence of chronic diseases including cancer and diabetes is increasing globally. Fruits and vegetables are potential sources of functional phytochemicals for prevention and management of these diseases. Bitter gourd, *Momordica charantia*, a popularly cultivated vegetable, contains a number of bioactive compounds with well-documented medicinal properties. These include cucurbitacin-B with proven anticancer activities, and charantin and plant insulin having demonstrated antidiabetic properties. However, the popularly grown varieties of bitter gourd are reportedly poor in the content of these phytomedicines and used only for vegetable purposes. We utilized a germplasm of wild bitter gourd, *M. charantia* var. *muricata* for the development of eight medicinal varieties including CBM3, CBM6, CBM10, CBM12, CBM13, CBM18, CBMH10 and CBMH12 in this crop, and evaluated the contents of these three phytomedicines in their fruits. We report here on the botanical description of these varieties, and phenotypic fruit traits and comparative performance of these varieties along with 17 popularly cultivated varieties with regard to the content of the three phytomedicines. We also report on the comparative performance with regard to fruit weight among the 25 varieties, and correlation of the contents of the three phytomedicines *inter se* and with fruit weight. Finally, we suggest five varieties including CBMH10, CBMH12, CBM12, CBM10 and CBM13, in that order, for consumption of their fruits as a source of functional foods for prevention and management of cancer and diabetes, and also as a source for extraction of these three phytomedicines for further studies on their pharmaceutical potential.

**KEYWORDS:** bitter gourd; breeding; medicinal crop-variety; anticancer phytomedicine; antidiabetic phytomedicine; functional food

---

### Open Access

Received: 23 Jan 2026

Accepted: 20 Apr 2026

Published: 24 Apr 2026

Copyright © 2026 by the author. Licensee Hapres, London, United Kingdom. This is an open access article distributed under the terms and conditions of Creative Commons Attribution 4.0 International License.

## INTRODUCTION

Prevalence of chronic diseases including cancer and diabetes is increasing globally. There were about 20 million new cases of cancer with 9.7 million deaths due to it in 2022 (<https://pressroom.cancer.org/GlobalCancerStatistics2024>). Similarly, 589 million adults were living with diabetes in 2024 and the number of deaths due to it was 3.4 million (<https://diabetesatlas.org/>). A number of crop plants under different commodity groups, specifically fruits and vegetables, are known to contain anticancer and antidiabetic bioactive compounds [1]. Hence, utilization of nutraceutical-rich crops, such as fruits and vegetables as a source of medicinally active phytochemicals is a potential alternative for prevention and management of these diseases [2].

Bitter gourd, also known as bitter melon or balsam pear, (*Momordica charantia* L.), is a potential cucurbit vegetable crop that can be popularized as a functional food crop. This plant contains over 60 phytomedicines active against over 30 diseases [3]. Several medicinal properties of bitter gourd including antidiabetic, anticancer, antitumor, antileukemic, antioxidant, antiulcer, anti-inflammatory, hypocholesterolemic and hypotriglyceridemic properties are well-documented in research [4]. Almost all parts of bitter gourd, primarily the fruits, contain cucurbitacin-B that have proven anticancer actions [5]. They also contain charantin and plant insulin (syn. p-insulin, polypeptide-p), that have clinically demonstrated hypoglycemic and antihyperglycemic activities and established beneficial effects on diabetes [6]. Plant insulin can control both type I and type II diabetes [7]. However, the popularly cultivated varieties of bitter gourd have lower content of phytomedicines [8] and are used only for vegetable purposes.

The traditional breeding programs in this crop aimed mainly at fruit yield along with quality, maturity time, fruit characteristics, economic traits, phytonutrients content etc. by utilizing local landraces and a number of varieties have been developed with improvement in these characters [9,10]. However, available literature does not evidence for any report on the development of varieties with enhanced content of phytomedicines in this crop. Assessment of bitter gourd germplasms has evidenced for sufficient genetic variation with regard to agromorphological and fruit-yield traits [11–13]. However, only limited germplasms exhibited sufficient genetic variability with regard to bioactive compounds required for genetic improvement [14–18].

The wild crop relatives of *Momordica* including *M. charantia* var. *muricata* (Willd.) Chakrav., *M. dioica* Roxb. ex Willd., *M. sahyadrica* Joseph and Antony, *M. subangulata* Kuroda and Habe, *M. cochinchinensis* (Lour.) Spreng., *M. balsamina* L., *M. cymbalaria* Hook. f., etc., possess wide genetic diversity for biotic stress resistance, abiotic stress tolerance, enhanced nutritional quality, high yielding traits, rare sex forms, unique fruit quality traits, etc. [19]. Among these, *M. charantia* var. *muricata* is considered as the progenitor of bitter gourd, *M. charantia* var. *charantia*, and is its closest

wild allied species [20,21]. We, therefore, explored a germplasm established through the collection from wildy grown bitter gourd plants belonging to the botanical variety *M. charantia* var. *muricata* (Willd.) Chakrav. (syn. *M. charantia* var. *abbreviata* Ser., *M. muricata* Willd.) for development of varieties with high contents of three anticancer and antidiabetic phytomedicines, mentioned earlier, and developed eight varieties, and report here. A compendium with comprehensive reviews on 50 crop plants belonging to diverse crop commodity groups provided the information on breeding for various nutraceuticals [1]. However, no crop variety was reported to be developed only as a source of phytomedicine, i.e., for ‘medicinal’ purposes. We have, therefore, performed breeding in wild bitter gourd with the sole objective to develop ‘medicinal crop-varieties’.

Since the weight of fruits is an important factor of total yield of bitter gourd [22] and thereby the total content of phytomedicines in each fruit, we studied the performance of these eight varieties along with 17 varieties grown popularly in six countries for ‘vegetable’ purposes with regard to both phytomedicine content and fruit weight, and also the nature and extent of correlation of the content of these three phytomedicines *inter se* and with fruit weight. This facilitated selection of the ideal varieties for consumption of their fruits and fruit products as a source of functional foods for prevention and management of cancer and diabetes and for use as a source for extraction of these phytomedicines for future pharmaceutical and bioavailability studies.

## MATERIALS AND METHODS

### Plant Materials

A germplasm of wildy grown plants of bitter gourd belonging to *Momordica charantia* var. *muricata* (Willd.) Chakrav., an allied botanical variety of the domesticated bitter gourd botanical variety, *M.c.* var. *charantia* L. was established by us previously at the Clemson University. Six plants from this germplasm were used for breeding of the first six medicinal varieties used in this study. Two of these medicinal varieties and a popularly grown variety with the heaviest fruits, Taiwan White, belonging to *M.c.* var. *charantia* L. were used for crossing to develop two hybrids. These eight varieties were grown in three consecutive times, called hereinafter as environments, under controlled conditions in a greenhouse following Complete Randomized Design with four replications along with 17 popularly grown vegetable varieties.

### **Recording of Botanical Descriptions and Fruit Phenotypic Traits**

Morphological characters on mature plants, leaves, flowers, fruits and seeds extracted from ripen fruits were recorded visually in the greenhouse. The quantitative data for leaves, flowers, fruits and seeds were recorded from the average of ten replicates of the corresponding plant parts.

The four fruit phenotypic traits including color, surface texture, luster and shape were recorded on visible observation in the greenhouse. The apparent size categories were derived arbitrarily from the average fruit weight of fruits as >1–25 g Very small, >25–50 g Small, >50–75 g Medium, >75–100 g Large and >100 g Very Large.

### **Recording of Data for Fruit Size**

The data for fresh fruit weight of the 25 varieties for the four replications were derived from average over the three environments and used for statistical analysis.

### **Extraction and Quantification of Phytomedicines**

Five fresh green fruits from each of the 25 varieties for the four replications in each environment were randomly selected and used for lyophilization. The freeze-dried powders of the three environments were pooled for each variety and each replication. The powders of the first two replications for each variety were pooled to derive the one replication for analysis. Similarly, the powders from the other two replications for that variety were pooled to derive another replication for analysis.

Quantification of the contents (mg/g) of one anticancer phytomedicine, cucurbitacin-B, and two antidiabetic phytomedicines, charantin and plant insulin, was done by HPLC analysis of the pooled lyophilized powders in two replications as described above and expressed in terms of mg/g of lyophilized powder. The detailed methods of extraction and quantification are described elsewhere [23].

### **Statistical Analysis**

Statistical analysis of variation in content of the phytomedicines was done by analysis of variance (ANOVA) following Completely Randomized Design (CRD) with two replications as mentioned above. Similar statistical analysis was done for fruit weight also but with four replications using averaged data over three environments as described earlier. The varietal effects were compared post-hoc using Duncan's Multiple Range Test (DMRT) and assigned alphabetical grouping ranks. Correlation analysis was done following routine statistical analysis.

## RESULTS

### Development of Medicinal Varieties and Their Hybrids

Pure-line selection until the 6th selfing generation ( $S_6$ ) from six wildy grown plants belonging to *Momordica charantia* var. *muricata* (Willd.) Chakrav. led to the development of six bitter gourd varieties including CBM3, CBM6, CBM10, CBM12, CBM13 and CBM18. Preliminary studies exhibited smaller fruit size and higher phytomedicine content of these varieties as compared to the popular vegetable cultivars. Two varieties, CBM10 and CBM12, selected from them based on higher phytomedicine content were hybridized, using them as male parents, with Taiwan White, a popularly cultivated vegetable variety with the largest fruit size belonging to *M. charantia* var. *charantia*, as the female parent, that resulted in two hybrid varieties, viz., CBMH10 and CBMH12. Results of evaluation of these eight varieties in comparison to the available vegetable varieties are presented in the following sections.

### Botanical Descriptions of the Varieties Developed

Several morphological features of the plants and their various parts of the eight medicinal varieties developed by us are presented in Table 1. Morphological features of the female parent of the two hybrid varieties, Taiwan White, are also furnished for comparative depiction. As expected, many qualitative features of the mature plants and their leaves, flowers, fruits and seeds were uniform among all the nine varieties. However, in this regard, the varieties exhibited different features for leaf color and seed surface, and this variation had no relation to the botanical varieties. Besides, the color of the dry seeds was black in Taiwan White and the two hybrids, while creamy in the CBM varieties. In general, the CBM varieties had less vigorous plants compared to Taiwan White, as enumerated by the quantitative attributes of the plants and their various parts. The two hybrids were intermediate in this regard as expected. Most importantly, this nature was more conspicuous for the fruit and seed attributes.

**Table 1.** Botanical descriptions of six medicinal varieties, female parent of the hybrids and two hybrid varieties of bitter gourd.


		CBM3	CBM6	CBM10	CBM12	CBM13	CBM18	Taiwan White	CBMH10	CBMH12
Botanical variety		<i>muricata</i>	<i>muricata</i>	<i>muricata</i>	<i>muricata</i>	<i>muricata</i>	<i>muricata</i>	<i>charantia</i>	<i>charantia</i> × <i>muricata</i>	<i>charantia</i> × <i>muricata</i>
Mature plants	Growth habit	Long vines	Long vines	Long vines	Long vines	Long vines	Long vines	Long vines	Long vines	Long vines
	Plant type	Pilose	Pilose	Pilose	Pilose	Pilose	Pilose	Pilose	Pilose	Pilose
	Diameter (mm)	13.0	15.0	14.0	9.0	17.3	13.0	26.7	17.0	16.3
Leaves	Blade shape	Ovate	Ovate	Ovate	Ovate	Ovate	Ovate	Ovate	Ovate	Ovate
	Blade form	Deep lobed	Deep lobed	Deep lobed	Deep lobed	Deep lobed	Deep lobed	Deep lobed	Deep lobed	Deep lobed
	Margin	Spiny	Spiny	Spiny	Spiny	Spiny	Spiny	Spiny	Spiny	Spiny
	Margin edges	Flat	Flat	Flat	Flat	Flat	Flat	Flat	Flat	Flat
	Width (cm)	10.8	12.4	10.0	8.3	10.5	11.2	18.0	12.2	17.1
	Length (cm)	8.9	8.9	8.5	6.4	8.5	7.1	18.0	10.7	15.7
	Leaf surface	Smooth	Smooth	Smooth	Smooth	Smooth	Smooth	Smooth	Smooth	Smooth
	Dorsal surface pubescence	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy
	Ventral surface pubescence	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy	Soft hairy
	Leaf color	Dark Green	Light green	Dark Green	Light green	Dark Green	Dark Green	Light green	Light green	Light green
	Petiole length (cm)	5.1	5.5	6.1	3.2	4.6	4.6	5.3	5.0	6.9
Flowers-Pistillate	Diameter (cm)	3.2	2.8	3.0	2.3	3.0	2.8	3.4	3.5	3.0
	Ovary	Fusiform	Fusiform	Fusiform	Fusiform	Fusiform	Fusiform	Fusiform	Fusiform	Fusiform
	Pedicel length	5.6	7.1	6.0	5.6	9.1	5.8	11.0	10.4	9.0
	Margin shape	Curved	Curved	Curved	Curved	Curved	Curved	Curved	Curved	Curved
	Margin edges	Plain	Plain	Plain	Plain	Plain	Plain	Plain	Plain	Plain
	Sepal width (mm)	1.2	1.0	1.2	1.0	1.1	1.1	2.0	1.3	1.3
	Sepal length (mm)	4.0	5.0	4.0	4.0	5.2	4.0	5.5	4.5	5.5
	Petal length (cm)	1.6	1.3	1.5	1.3	1.5	1.4	1.6	1.6	1.7
	Petal width (cm)	1.1	0.9	0.9	0.9	1.4	0.9	1.7	1.5	1.2
Flowers-Staminate	Diameter (cm)	3.8	3.4	3.1	3.1	3.8	3.2	4.6	4.3	3.5
	Sepal length (mm)	6.3	6.5	6.0	6.0	6.5	6.0	9.0	7.0	4.5
	Sepal width (mm)	3.3	5.0	3.0	4.0	4.0	3.6	5.0	5.0	4.0
	Petal length (cm)	1.9	1.7	1.6	1.6	2.1	1.6	4.1	2.0	1.6
	Petal width (cm)	1.2	1.1	0.9	0.9	1.2	0.9	4.0	1.7	1.4
	Pedicel length (cm)	5.4	6.0	3.5	5.1	5.8	4.4	7.4	6.9	7.2
	Color	Lemon yellow	Lemon yellow	Lemon yellow	Lemon yellow	Lemon yellow	Lemon yellow	Lemon yellow	Lemon yellow	Lemon yellow
Fruits	Length (cm)	10.8	8.7	6.8	4.9	8.6	8.8	14.5	12.9	8.8
	Middle width (cm)	3.6	3.5	3.1	2.9	3.4	2.8	5.7	4.1	4.3
	Average weight (g)	41.7	33.1	4.9	39	30.2	5.9	146.3	64.5	74.4
Fruit stalks	Length (cm)	6.5	8.0	8.0	5.7	8.0	6.9	12.0	7.5	7.0
	Width (cm)	1.0	1.0	1.1	1.0	1.0	1.0	1.6	1.1	0.9
Seeds	Surface	Spiny	Spiny	Non-spiny	Spiny	Non-spiny	Non-spiny	Non-spiny	Non-spiny	Non-spiny
	Dry color	Creamy	Creamy	Dark creamy	Creamy	Creamy	Creamy	Black	Black	Black
	Length (mm)	12.8	13.5	13.3	12.3	13.8	13.8	13.8	14.0	13.2
	Width (mm)	6.5	6.5	6.5	7.2	7.5	6.8	7.8	7.8	7.5
	Thickness (mm)	3.5	3.3	3.2	3.5	3.8	3.3	3.8	3.8	3.5
	Weight/100 seeds (g)	15.0	17.0	16.2	16.5	18.2	16.3	19.5	18.6	17.7
	No. of seeds/ fruit	11.0	7.0	5.0	3.0	4.0	4.0	14.0	19.0	10.0

### Fruit Phenotypic Traits of the Varieties

Phenotypic traits of five fruit characters including color, surface texture, luster, shape, and apparent size of the eight medicinal varieties and 17 vegetable varieties grown popularly in six Asian countries are presented in Table 2. All the six CBM varieties and the two hybrids bore green, spiny and dull fruits. The nine popular vegetable varieties belonging to the same botanical variety, *muricata*, had also green, spiny and dull fruits. On the other hand, the eight popular vegetable varieties belonging to the *charantia* botanical variety had fruits of varied color and luster but all being non-spiny. One of these *charantia* varieties, viz. Large Top, had dull fruits unlike the other seven varieties bearing glossy fruits. The three Taiwanese varieties, viz. Hybrid White Pearl, Hybrid Taiwan White and Taiwan White, had white fruits, the remaining five *charantia* varieties having green fruits. Fruit shape was, in general, long except the four *muricata* varieties, viz. CBM12, Hybrid India Baby, Small Baby and Hybrid Baby Doll, and one hybrid variety, viz. CBMH12. All the six CBM varieties had very small- and small-sized fruits whereas the nine vegetable varieties belonging also to the *muricata* botanical variety bore very small, small, medium and large fruits. This trait was quantified and is elaborated in the Section, Variation in Fruit Size. The *charantia* vegetable varieties had either medium or large fruits but one of them, viz., Taiwan White, bore very large fruits. The two hybrids, obviously, bore medium-sized fruits. These five contrasting phenotypic fruit traits are depicted in Figure 1.

**Table 2.** Fruit phenotypes of the 25 varieties of bitter gourd.

Varieties	Origin	Colour	Surface texture	Luster	Shape	Apparent Size
<b><i>M.c. var. muricata</i></b>						
CBM3	USA	Green	Spiny	Dull	Long	Small
CBM6	USA	Green	Spiny	Dull	Long	Small
CBM10	USA	Green	Spiny	Dull	Long	Very small
CBM12	USA	Green	Spiny	Dull	Round	Very small
CBM13	USA	Green	Spiny	Dull	Long	Small
CBM18	USA	Green	Spiny	Dull	Long	Very small
Hybrid India Star	India	Green	Spiny	Dull	Long	Medium
Hybrid India Baby	India	Green	Spiny	Dull	Round	Very small
Hybrid India Pearl	India	Green	Spiny	Dull	Long	Medium
India Long Green	India	Green	Spiny	Dull	Long	Very small
Hybrid India Green Queen	India	Green	Spiny	Dull	Long	Small
Japan Green Spindle	Japan	Green	Spiny	Dull	Long	Very small
Japan Long	Japan	Green	Spiny	Dull	Long	Large
Small Baby	Thailand	Green	Spiny	Dull	Round	Very small
Hybrid Baby Doll	Thailand	Green	Spiny	Dull	Round	Very small
<b><i>M.c. var. charantia</i></b>						
Hybrid Beauty Winner	China	Green	Non-spiny	Glossy	Long	Medium
Large Top	China	Green	Non-spiny	Dull	Long	Large
Hong Kong Green	Hong Kong	Green	Non-spiny	Glossy	Long	Medium
Hybrid Jumbo	Thailand	Green	Non-spiny	Glossy	Long	Medium
Hybrid Bangkok Large	Thailand	Green	Non-spiny	Glossy	Long	Large
Hybrid White Pearl	Taiwan	White	Non-spiny	Glossy	Long	Large
Hybrid Taiwan White	Taiwan	White	Non-spiny	Glossy	Long	Medium
Taiwan White	Taiwan	White	Non-spiny	Glossy	Long	Very Large
<b><i>M.c. var. charantia</i> × <i>M.c. var. muricata</i> hybrids</b>						
CBMH10	USA	Green	Spiny	Dull	Long	Medium
CBMH12	USA	Green	Spiny	Dull	Round	Medium



**Figure 1.** Contrasting fruit traits observed in the 25 varieties of bitter melon depicted with four representative varieties. (a) The parent varieties, CBM10 and Taiwan White, and their hybrid CBMH10. (b) The parent varieties, CBM12 and Taiwan White, and their hybrid CBMH12.

### Variation in Phytomedicine Content in Fruits

Statistical analysis of the contents of the three phytomedicines as mg/g of lyophilized powers of fresh fruits of the 25 varieties including the eight CBM varieties and 17 popular vegetable cultivars revealed highly significant variation (Table 3). The contents of cucurbitacin-B were significantly the highest in CBM10 (0.645 mg/g) and CBMH10 (0.615 mg) followed by CBM13 (0.5 mg). CBM18 also had considerable (0.465 mg) cucurbitacin-B content. Two popular vegetable varieties, Hybrid White Pearl and Taiwan White were equally the lowest performing varieties containing 0.12 mg and 0.13 mg, respectively. With respect to charantin, CBMH10 (8.0 mg) again exhibited significantly the highest content closely followed by CBM12 (7.88 mg). One hybrid variety, CBMH12, two CBM varieties, CBM10 and CBM3 along with four popular hybrid vegetable varieties, viz., Hybrid Beauti Winner, Hybrid Jumbo, Hybrid White Pearl and Hybrid India Green Queen had considerable charantin contents. Hybrid India Star (4.2 mg) followed by Hybrid Baby Doll (5.37 mg) had the lowest charantin content. CBMH10 (4.0 mg) had the highest content of

plant insulin and was seconded by CBMH12 (0.3 mg) that was closely followed by CBM12 and CBM3 with similar (0.26 mg) content. Four CBM varieties including CBM13 and CBM18 followed by CBM10 and CBM6 had also considerable charantin contents ranging between 0.185 to 0.210 mg. Six cultivars including Hybrid Beauty Winner, Hong Kong Green, Taiwan White, Hybrid Bangkok Large, Hybrid India Star and Hybrid Taiwan White had equally the least (0.11–0.12 mg) plant insulin contents.

**Table 3.** Mean contents (mg/g) of cucurbitacin-B (CCR), charantin (CHR) and plant insulin (PLIN) in lyophilized powder of fresh fruits and mean fresh fruit weight (FR WT) in 25 bitter gourd varieties over three replications.

Genotype	CCR	Rank <sup>†</sup>	CHR	Rank	PLIN	Rank	FR WT	Rank
<b><i>M.c. var. muricata</i></b>								
CBM3	0.360	cdefg	7.470	abc	0.260	bc	41.67	e
CBM6	0.420	bcde	6.455	cdef	0.185	de	33.13	ef
CBM10	0.645	a	7.580	abc	0.190	de	4.90	h
CBM12	0.435	bcd	7.880	ab	0.260	bc	3.84	h
CBM13	0.500	b	6.590	bcdef	0.210	cd	30.20	ef
CBM18	0.465	bc	7.305	abcd	0.205	cd	5.89	h
Hybrid India Star	0.150	ijk	4.200	g	0.110	f	74.83	bcd
Hybrid India Baby	0.400	bcdef	7.100	abcde	0.160	def	4.28	h
Hybrid India Pearl	0.210	hijk	6.950	abcde	0.160	def	72.26	bcd
India Long Green	0.330	defgh	6.420	cdef	0.140	ef	22.64	fg
Hybrid India Green Queen	0.300	efgh	7.435	abc	0.140	ef	25.29	fg
Japan Green Spindle	0.350	cdefg	6.950	abcde	0.140	ef	16.10	gh
Japan Long	0.240	ghijk	7.270	abcd	0.170	def	78.69	bc
Small Baby	0.410	bcdef	6.420	cdef	0.130	ef	3.48	h
Hybrid Baby Doll	0.290	fgh	5.370	f	0.130	ef	5.10	h
<b><i>M.c. var. charantia</i></b>								
Hybrid Beauty Winner	0.150	ijk	7.480	abc	0.120	f	74.81	bcd
Large Top	0.270	ghi	7.160	abcde	0.130	ef	83.87	b
Hong Kong Green	0.150	ijk	7.120	abcde	0.115	f	70.61	cd
Hybrid Jumbo	0.260	ghij	7.460	abc	0.130	ef	70.87	cd
Hybrid Bangkok Large	0.150	ijk	7.260	abcd	0.110	f	78.50	bc
Hybrid White Pearl	0.120	k	7.440	abc	0.170	def	82.37	bc
Hybrid Taiwan White	0.140	jk	5.850	ef	0.110	f	63.600	d
Taiwan White	0.130	k	5.985	def	0.115	f	146.33	a
<b><i>M.c. var. charantia</i> × <i>M.c. var. muricata</i> hybrids</b>								
CBMH10	0.615	a	8.000	a	0.400	a	64.47	d
CBMH12	0.365	cdefg	7.600	abc	0.300	b	74.36	bcd
F-Value	15.21		4.82		15.76			
p-value	$1.15 \times 10^{-9}$ ***		$1.07 \times 10^{-4}$ ***		$7.72 \times 10^{-10}$ ***			
SE(m)	0.038		0.389		0.018			
CV%	17.27		7.98		14.56			

<sup>†</sup> Alphabetical grouping rank derived from Duncan's Multiple Range Test. \*\*\* indicates statistically highly significant at  $p < 0.001$ .

### Variation in Fruit Size

Fruit weight also exhibited highly significant variation among the varieties (Table 3). The varieties belonging to the *charantia* varieties bore, in general, larger fruits weighing between 63.6 g to 146.33 g. The *muricata* varieties had varied sizes ranging between 3.84 g to 78.69 g. The six CBM varieties bore smaller fruits weighing between 3.84 g to 41.67 g.

Statistically six varieties including CBM12 (3.84 g), CBM10 (4.9 g) and CBM18 (5.89 g) had the smallest fruits whereas Taiwan White (146.33 g) had by far the largest fruits among all. As expected, the two hybrid varieties, CBMH12 (74.36 g) and CBMH10 (64.47 g) had medium sized fruits (Figure 1).

#### Associations among Phytomedicine Contents and Fruit Weight

Correlation analysis among the contents of three phytomedicines and fruit weight revealed highly significant positive association ( $r = 0.672$ ) between cucurbitacin-B content and plant insulin content. Charantin content and plant insulin content were also highly and positively correlated ( $r = 0.518$ ). Interestingly, fruit weight had highly significant negative association with cucurbitacin-B ( $r = -0.65$ ). Correlations of other combinations were statistically non-significant (Table 4).

**Table 4.** Correlation of contents of cucurbitacin-B (CCR-B), charantin (CHR) and plant insulin (PLIN) *inter se* and with fruit weight (Fr Wt) among 25 bitter gourd varieties.

		CCR-B	CHR	PLIN	Fr Wt
CCR-B	Pearson Correlation	1	0.362	0.672 **	-0.650 **
	Sig. (2-tailed)		0.076	0.000	0.000
	N	25	25	25	25
CHR	Pearson Correlation	0.362	1	0.518 **	-0.083
	Sig. (2-tailed)	0.076		0.008	0.695
	N	25	25	25	25
PLIN	Pearson Correlation	0.672 **	0.518 **	1	-0.150
	Sig. (2-tailed)	0.000	0.008		0.473
	N	25	25	25	25
Fr Wt	Pearson Correlation	-0.650 **	-0.083	-0.150	1
	Sig. (2-tailed)	0.000	0.695	0.473	
	N	25	25	25	25

\*\* Correlation is significant at the 0.01 level (2-tailed).

#### DISCUSSION

Several crop plants contain therapeutically and nutritionally important phytochemicals. A recently published compendium deliberated on the biochemistry and medicinal importance of these health-related nutraceuticals in 50 different crop plants [1]. It has also presented reviews on traditional breeding, transgenics and molecular breeding for genetic improvement in these crops aiming at the improvement in usual agronomic or horticultural traits such as yield, quality, stress resistance and also content of bioactive phytochemicals with them in a few cases. Similarly, previous breeding programs in bitter gourd, a popular vegetable crop, aimed only at horticultural traits, primarily on fruit yield and stress resistance, and produced varieties to be used for vegetable purposes [24]. On the other hand, our breeding program in bitter gourd was aimed only at higher content of medicinally important bioactive compounds and to

develop medicinal crop-varieties. The earlier bitter gourd varieties were developed by using pure-line selection and hybridization [4,9,24]. However, it is not clear whether any of their parent materials belonged to *M.c. var. muricata*.

*M.c. var. muricata* (syn. *M.c. var. abbreviata* ser.) is considered to be more potent in disease prevention than its cultivated form [25]. We explored, therefore, a germplasm of wild bitter gourd in our breeding program to develop medicinal varieties. Botanical descriptions of these varieties were as reported earlier specifically having small-sized and spiny fruits and smaller seed size [26–28]. With regard to the phenotypic traits of fruits, the CBM varieties were uniformly green, spiny, dull and small-sized. The two hybrids derived from them were also green, spiny and dull evidencing for dominance of these three traits over white, non-spiny and glossy and supporting the observations of Kole et al. (2012) [29]. They also observed difference in seed color between CBM12 and Taiwan White as observed by us (Table 1). The medium-sized fruits in the two hybrids observed by us substantiate the quantitative nature of fruit weight as observed earlier [29]. However, further genetic studies are required to detect the nature of inheritance of fruit shape although the observation in our work indicates possible dominance of round over long size as the hybrid of CBM12 with round fruits and Taiwan White with long fruits had round fruits. Crossability between *M.c. var. charantia* and *M.c. var. muricata* and characterization of their interspecific hybrids based on morphology, pollen-pistil compatibility, cytology, palynology, molecular relation among the parents and hybrids were studied earlier [20,30,31]. But these studies were not meant for detection of the nature of inheritance of economically important traits or development of interspecific hybrid varieties. A recent study reported significant heterosis with regard to charantin content in F<sub>1</sub> hybrids of three crosses involving some bitter gourd lines [32]. However, available literature does not provide report on the development of any hybrid variety with enhanced charantin or any other bioactive compounds from the above or any other cross(es).

The eight bitter gourd varieties of the CBM series had, in general, higher contents of all the three anticancer and antidiabetic phytochemicals in comparison to the 17 popularly grown vegetable varieties belonging to both *muricata* and *charantia* botanical varieties. Among them, CBMH10 was found to be universally the richest. CBM10 also assumed the first position along with CBMH10 for cucurbitacin-B content and the third position with regard to charantin content. CBM12 ranked second for charantin content and third for plant insulin content while the hybrid derived by using it ranked third for charantin and second for plant insulin. Another CBM variety, viz., CBM13, acquired the second rank with regard to cucurbitacin-B content. None of the remaining 20 varieties ranked first and/or second for any of the three phytochemicals. Therefore, consumption of fresh fruits of the above five varieties and their extracts or powders as functional foods may help in mitigation of the problems of

cancer and diabetes. Since diabetes leads also to blindness, kidney failure, heart attacks, stroke and lower limb amputation (<https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>), use of these varieties as functional foods may help in controlling these diseases. In addition, these varieties may be used as sources for extraction of the three phytomedicines for further pharmaceutical and bioavailability studies with emphasis on their potential use for mitigation of cancer and diabetes. Studies on other phytomedicines in these varieties as reported to be present in bitter gourd [3] and their potential medicinal utilities [4] may also be taken up in future.

Fruit weight is an important factor of fruit yield [22] and thereby total phytomedicine contained in fruits. The CBM varieties had, in general, smaller fruits as compared to the popular cultivars. However, the two hybrids CBMH12 and CBMH10 bore heavier fruits equaling to many popular cultivars. These two hybrids, therefore, emerged as superior to all the varieties considering both medicinal and vegetable purposes and may be recommended as 'dual-purpose' varieties.

The correlation study has led to an interesting observation of negative association between fruit weight and cucurbitacin-B content indicating the requirement of compromising fruit yield for utilization of bitter gourd varieties as a source of this anticancer phytomedicine. Another interesting observation was the positive correlation of the contents of the two antidiabetic phytomedicines, charantin and plant insulin suggesting future studies on genetic and biochemical mechanisms underlying their production.

All the CBM varieties were developed by either selection or hybridization, both traditional breeding methods, and so their fruits and fruit products can be safely consumed and these varieties may enjoy acceptance by both the farmers and consumers.

## CONCLUSIONS

The results from this study demonstrates the importance of developing crop varieties with higher contents of therapeutic bioactives for their use as sources of functional foods aiming at prevention and management of human diseases including cancer and diabetes. They also evidenced that wild crop relatives could be a potential genetic resource for developing medicinal varieties in crops.

## DATA AVAILABILITY

The detailed data are not publicly available but we will be happy to share data on reasonable request.

### AUTHOR CONTRIBUTIONS

Conceptualization, CK; methodology, CK and PK; validation, CK and PK; formal analysis, CK and PK; investigation, CK and PK; data curation, PK; writing—original draft preparation, CK; writing—review and editing, CK and PK; supervision, CK. Both authors have read and agreed to the published version of the manuscript.

### CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

### FUNDING

The first author acknowledges USDA National Institute of Food and Agriculture for financial support through the Grant No. 2009-35301-15041 for the original work. The authors declare that no funds, grants, or other support were received during the final analysis, interpretation and preparation of this manuscript.

### ACKNOWLEDGMENTS

The authors acknowledge Albert G. Abbott of the Department of Genetics and Biochemistry and Richard K. Marcus of the Department of Chemistry, Clemson University, Clemson, SC, USA for laboratory facilities; Bode Olukolu, while working at the Department of Genetics and Biochemistry, Clemson, SC, USA for assistance in extraction and quantification of phytomedicines; and Deb Sankar Gupta from the Uttar Banga Krishi Viswavidyalaya, India for assistance in statistical analysis.

### REFERENCES

1. Kole C, editor, Compendium of crop genome designing for nutraceuticals. Nature MRW Plant Sciences. Cham (Switzerland): Springer Nature. 2024.
2. Mahmoud S, Ibrahim AAA. Fruits and vegetables as sources of functional phytochemicals for the prevention and management of obesity, diabetes, and cancer. In: Egbuna C, Hassan S, editors, Dietary phytochemicals. Cham (Switzerland): Springer Nature; 2021. pp. 147-67. doi: 10.1007/978-3-030-72999-8\_8.
3. Raman A, Lau C. Anti-diabetic properties and phytochemistry of *Momordica charantia* L. (Cucurbitaceae). Phytomedicine. 1996;2(4):349-62. doi: 10.1016/S0944-7113(96)80080-8
4. Behera TK, Matsumura H, Kole C. Glimpse on genomics and breeding in bitter gourd: a crop of the future for food, nutrition and health security. In: Kole C, Matsumura H, Behera TK, editors, The bitter gourd genome. Cham (Switzerland): Springer Nature; 2020, pp. 1-6. doi: 10.1007/978-3-030-15062-4\_1

5. Nie W, Wang Y, Tian X, Liu J, Jin Z, Xu J, et al. Cucurbitacin B and its derivatives: A review of progress in biological activities. *Molecules*. 2024;29(17):4193. doi: 10.3390/molecules29174193
6. Joseph B, Jini D. Antidiabetic effects of *Momordica charantia* (bitter melon) and its medicinal potency. *Asian Pac J Trop Dis*. 2013;3(2):93-102. doi: 10.1016/S2222-1808(13)60052-3
7. Chakraborty P. Development of new anti-diabetic drug from medicinal plant-genomic research. *J Diabetol*. 2018;2(1):6-7.
8. Kole C, Olukolu BA, Kole P, Abbott AG. Association mapping of fruit traits and phytomedicine contents in a structured population of bitter melon (*Momordica charantia* L.). In: Thies JA, Kousik S, Levi A, editors, *Cucurbitaceae 2010 proceedings*. Alexandria (VA, US): American Society for Horticultural Science; 2010, pp. 42-45.
9. Behera TK, Bhardwaj DR, Gautam KK. Bitter gourd: breeding and genomics. *Veg Sci*. 2023;50(Special Issue):189-207. doi: 10.61180/vegsci.2023.v50.spl.06
10. Rao PG. Recent advances in breeding of bitter gourd (*Momordica charantia* L.). In: Al-Khayri JM, Jain SM, Johnson DV, editors, *Advances in plant breeding strategies: vegetable crops*. Cham (Switzerland): Springer Nature; 2021; pp. 87-121. doi: 10.1007/978-3-030-66961-4\_3
11. Alhariri A, Behera TK, Jat GS, Devi MB, Boopalakrishnan G, Teleb AA et al. Analysis of genetic diversity and population structure in bitter gourd (*Momordica charantia* L.) using morphological and SSR markers. *Plants*. 2021;10(9):1860. doi: 10.3390/plants10091860
12. Bhati S, Prakash S, Kumar V, Singh B, Singh MK, Gangwar LK et al. Genetic variability, heritability and genetic advance for yield and yield contributing characters in bitter gourd (*Momordica charantia* L.) genotypes. *Int J Environ Clim*. 2023;13:1867-75. doi: 10.9734/ijecc/2023/v13i92418
13. Mallikarjuna KN, Tomar BS, Mangal M, Singh N, Singh D, Kumar S et al. Genetic diversity and population structure analyses in bitter gourd (*Momordica charantia* L.) based on agro-morphological and microsatellite markers. *Plants*. 2023;12(19):3512. doi: 10.3390/plants12193512
14. Hazra P, Hazra S, Acharya B, Dutta S, Saha S, Mahapatra P et al. Diversity of nutrient and nutraceutical contents in the fruits and its relationship to morphological traits in bitter gourd (*Momordica charantia* L.). *Sci Hortic*. 2022;305:111414. doi: 10.1016/j.scienta.2022.111414
15. Kim AK, Lee HJ, Oh MM, Lee WM, Lee SG, Chae WB et al. Selection of bitter gourd (*Momordica charantia* L.) germplasm for improvement anti-diabetic compound contents. *Korean J Breed Sci*. 2013;45(4):332-8. doi: 10.9787/KJBS.2013.45.4.332
16. Mahwish M, Saeed F, Nisa MU, Nadeem MT. Minerals and phytochemical analysis of bitter melon fruits and its components in some indigenous and exotic cultivars. *Bioscience J*. 2018;34(6):1622-31. doi: 10.14393/BJ-v34n6a2018-39827

17. Perez JL, Jayaprakashs GK, Crosby K, Patil BS. Evaluation of bitter melon (*Momordica charantia*) cultivars grown in Texas and levels of various phytonutrients. *J Sci Food Agric*. 2019;99(1):379-90. doi: 10.1002/jsfa.9199
18. Zhou S, Allard PM, Wolfrum C, Ke C, Tang C, Ye Y et al. Identification of chemotypes in bitter melon by metabolomics: a plant with potential benefit for management of diabetes in traditional Chinese medicine. *Metabolomics*. 2019;15(8):104. doi: 10.1007/s11306-019-1565-7
19. Devi S, Neelavathi R, Indu Rani C, Kalaiyarasi R, Gurusamy K. Utilization of crop wild relatives in genetic improvement of bitter gourd, *Momordica charantia* L. *Genet Resour Crop Evol*. 2026;73(4):170. doi: 10.1007/s10722-026-02768-x
20. Rathod V, Behera TK, Das Munshi A, Vinod, Jat GS. Crossability studies among *Momordica charantia* var. *charantia* and *Momordica charantia* var. *muricata*. *Indian J Agric Sci*. 2019;89(11):1900-5. doi: 10.56093/ijas.v89i11.95339
21. Bano M, Sharma G. *Momordica charantia* var. *Muricata* L.: need to conserve and promote its cultivation in Jammu province (J&K). *Int J Pharm Biol Sci*. 2017;8(4):518-21. doi: 10.22376/ijpbs.2017.8.4.b518-521
22. Reddy MS, Prasanth P, Laxminarayana D, Saidaiah P. Correlation and path coefficient analysis of fruit yield and yield attributes in twenty seven genotypes of bitter gourd (*Momordica Charantia* L.). *Pharma Innov J*. 2022;11(10):1383-89.
23. Kole C, Kole P, Randunu KM, Choudhary P, Podila R, Ke PC, et al. Nanobiotechnology can boost crop production and quality: first evidence from increased plant biomass, fruit yield and phytomedicine content in bitter melon (*Momordica charantia*). *BMC Biotechnol*. 2013;13:37. doi: 10.1186/1472-6750-13-37
24. Behera TK, Behera S, Bharathi LK, Joseph JK. Bitter gourd: botany, horticulture and breeding. In: Janick J, editor, *Horticulture reviews*. Hoboken (NJ, US): Wiley, Blackwell. 2020; p 101-41.
25. Lii C-K, Chen H-W, Yun W-T, Liu KL. Suppressive effects of wild bitter gourd (*Momordica charantia* Linn. var. *abbreviata* ser.) fruit extracts on inflammatory responses in RAW 264.7 macrophages. *J Ethnopharmacol*. 2009;122(2):227-33. doi: 10.1016/j.jep.2009.01.028
26. Asna AC, Joseph J, John KJ. Botanical description of bitter gourd. In: Kole C, Matsumura H, Behera TK, editors, *The bitter gourd genome*. Cham (Switzerland): Springer Nature. 2020, p 7-31. doi: 10.1007/978-3-030-15062-4\_2
27. Suma A, Alfia MA, John KJ, Pradheep K, Harish GD, Thirumalaisamy PP, et al. On the development of descriptors in small bitter gourd (*Momordica charantia* L. var. *muricata* (Willd.) Chakrav). *Genet Resour Crop Evol*. 2023;70:289-308. doi: 10.1007/s10722-022-01494-4
28. Sheemole MS, Antony VT. Study of antioxidant activity and phytochemical analysis of fruits of *Momordica charantia* var. *muricata* using LC-MS. *World J Biol Pharm Health Sci*. 2025;21(03):558-64. doi: 10.30574/wjpbphs.2025.21.3.0313

29. Kole C, Olukolu BA, Kole P, Rao VK, Bajpai A, Backiyarani S, et al. The first genetic map and positions of major fruit trait loci of bitter melon (*Momordica charantia*). J Plant Sci Mol Breed. 2012;1(1):1-6. doi: 10.7243/2050-2389-1-1
30. Haseena Bai N, Suhara Beevy S. Characterization of intraspecific F<sub>1</sub> hybrids of *Momordica charantia* L. based on morphology, cytology and palynology. Cytologia. 2012;77(3):301-10. doi: 10.1508/cytologia.77.301
31. Bharathi LK, Munshi AD, Behera TK, Vinod, John KJ, Das AB, et al. Production and preliminary characterization of inter-specific hybrids derived from *Momordica* species. Curr Sci. 2012;103(2):178-86.
32. Srinivasulu B, Vijaya Padma SS, Naram Naidu L, Kumar CK, Sekhar V. Development of hybrids in bitter gourd for charantin and other qualitative traits. J Appl Hortic. 2024;26(1):78-82. doi: 10.37855/jah.2024.v26i01.15

How to cite this article:

Kole C, Kole P. Development of the First Medicinal Crop-Varieties with Enhanced Bioactive Compounds from Bitter Gourd (*Momordica charantia* L.). Crop Breed Genet Genom. 2026;8(2):e260012. <https://doi.org/10.20900/cbagg20260012>.